









CHEMISTRY (043) MARKING SCHEME 2016

SET-56/1/C

| Q | VALUES POINTS | MARKS |
|---|--|----------------|
| 1 |  | 1 |
| 2 | NO ₂ | 1 |
| 3 | (i) Molecular Solid - I ₂ (ii) Ionic Solid - NaCl <div style="text-align: right;">(Any other suitable example)</div> | ½ + ½ |
| 4 | 2- Phenylethanol | 1 |
| 5 | Like charged particles cause repulsion / Brownian movement / solvation | 1 |
| 6 | (i) Gas B , Higher the value of K_H lower is the solubility of gas / $p = K_H \cdot x$ (ii) Negative deviation from Raoult's law | ½ + ½ 1 |
| 7 | (i)  ii)  | 1+1 |
| | OR | |
| 7 | (i) $2\text{Fe}^{3+} + \text{SO}_2 + 2\text{H}_2\text{O} \longrightarrow 2\text{Fe}^{2+} + \text{SO}_4^{2-} + 4\text{H}^+$ (ii) $\text{XeF}_4 + \text{SbF}_5 \longrightarrow [\text{XeF}_3]^+ [\text{SbF}_6]^-$ | 1 1 |
| 8 | (i) [Co (NH ₃) ₆] Cl ₃ (ii) Hexaamminecobalt(III) chloride | 1 1 |
| 9 | (i) Zero order reaction, Molecularity is 2 / bimolecular reaction (ii) mol L ⁻¹ s ⁻¹ | ½ + ½ 1 |

| | | |
|----|---|------------------|
| | $= \frac{2.303}{300 \text{ s}} \times 0.4771$ $= \frac{1.099}{300 \text{ s}}$ $= 0.0036 \text{ s}^{-1} \quad / \quad 3.66 \times 10^{-3} \text{ s}^{-1}$ <p style="text-align: right;">(deduct ½ mark if unit is not written)</p> | 1 |
| 13 | <p>i) Liquid loving/ solvent loving.</p> <p>ii) Potential difference between the fixed layer and diffused / double layer of opposite charges</p> <p>iii) Some substances at higher concentration exhibit colloidal behaviour due to formation of aggregates. The aggregated particles thus formed are called associated colloids or micelles</p> | 1 1 1 |
| 14 | <p>(i) Mond's Process</p> <p>(ii) The melting point of alumina is very high. It is dissolved in cryolite which lowers the melting point and brings conductivity / acts as a solvent.</p> <p>(iii) Limestone is decomposed to CaO ,which removes silica impurity of the ore as slag.</p> $\begin{array}{ccc} \text{CaCO}_3 & \xrightarrow[\text{OR}]{\Delta} & \text{CaO} + \text{CO}_2 \\ \text{CaO} + \text{SiO}_2 & \longrightarrow & \text{CaSiO}_3 \\ & & \text{Slag} \end{array}$ | 1 1 1 |
| 15 | $\Delta T_b = i K_b \cdot m$ $i=2$ $= i \times K_b \times \frac{w_2 \times 1000}{M \times W_1}$ $= 2 \times 0.52 \text{ K kg mol}^{-1} \times \frac{4 \text{ g} \times 1000 \text{ g / kg}}{120 \text{ g/mol} \times 100 \text{ g}}$ $= \frac{2 \times 0.52}{3}$ $= 0.346 \text{ K}$ <p>Boiling point of water = 373.15 K / 373 K</p> $T_b = T_b^{\circ} + \Delta T_b$ $= 373.15 \text{ K} + 0.346 \text{ K} \quad / \quad 373 \text{ K} + 0.346 \text{ K}$ $= 373.496 \text{ K} \quad / \quad 373.346 \text{ K}$ | ½ 1 ½ 1 |
| 16 | <p>i) Because stability of higher oxidation state decreases as we move down the group / S is more stable in higher (+6) oxidation state whereas Te is more stable in +4 oxidation state.</p> <p>(ii) Due to absence of d orbital.</p> | 1 1 |

| | | |
|----|---|----------------------------|
| | (iii) Because I – Cl bond is weaker than I-I bond. | 1 |
| 17 | <p>(a)</p> $\text{CH}_3\text{OH} + \text{CH}_3 - \underset{\text{CH}_3}{\overset{\text{CH}_3}{\text{C}}} - \text{I}$ <p>(b)</p> $\text{CH}_3\text{CH}_2 - \underset{\text{O}}{\overset{\parallel}{\text{C}}} - \text{CH}_3$ <p>(c)</p>  | <p>1</p> <p>1</p> <p>1</p> |
| 18 | <p>(i) Aniline is a Lewis base while AlCl_3 is lewis acid. They combine to form a salt.</p> <p>(ii) Due to combined + I and solvation effects.</p> <p>(iii) Due to presence of H-bonding in primary amines.</p> | <p>1</p> <p>1</p> <p>1</p> |
| 19 | <p>(i)</p> $2 \text{ } \text{C}_6\text{H}_5\text{Cl} + 2\text{Na} \xrightarrow[\text{Ether}]{\text{dry}} \text{C}_6\text{H}_5\text{C}_6\text{H}_5 + 2\text{NaCl}$ <p>(ii) $\text{CH}_3\text{CH}=\text{CH}_2 \xrightarrow{\text{HBr / peroxide}} \text{CH}_3\text{CH}_2\text{CH}_2\text{Br} \xrightarrow{\text{NaI/acetone}} \text{CH}_3\text{CH}_2\text{CH}_2\text{I}$</p> <p>(iii)</p> $\text{CH}_3\text{CH}_2\underset{\text{Br}}{\text{CH}}\text{CH}_3 \xrightarrow{\text{Alc.KOH}} \text{CH}_3\text{CH}=\text{CHCH}_3$ | <p>1</p> <p>1</p> <p>1</p> |
| | | |

| | | |
|----|---|---|
| | OR | |
| 19 | <p>(i)</p>  <p>(ii)</p>  <p>(iii) $\text{CH}_3\text{CH}_2\text{NC}$</p> | <p>1</p> <p>1</p> <p>1</p> |
| 20 | <p>(i) On vulcanization, sulphur forms cross links at the reactive sites of double bond, the rubber gets stiffened.</p> <p>(ii) Ethylene glycol / $\text{HO}-\text{CH}_2\text{CH}_2-\text{OH}$, Terephthalic acid / </p> <p>(iii) Neoprene < Polythene < Terylene</p> | <p>1</p> <p>1</p> <p>1</p> |
| 21 | <p>(i) Starch - Polymer of α-D- glucose units / Polymer of α- glucose units. Cellulose - polymer of β-D- glucose units / polymer of β-glucose units.</p> <p>(ii) Phosphodiester linkage</p> <p>(iii) Fibrous protein - Keratin / myosin / collagen Globular protein - haemoglobin / insulin</p> | <p>1</p> <p>1</p> <p>$\frac{1}{2} + \frac{1}{2}$</p> |
| 22 | <p>(i) sp^3d^2, paramagnetic, high spin</p> <p>(ii)</p>  | <p>$1 + \frac{1}{2} + \frac{1}{2}$</p> <p>1</p> |
| 23 | (i) Caring nature, supportive, aware (or any other two suitable values) | $\frac{1}{2} + \frac{1}{2}$ |

| | | |
|----|--|---------------------------------------|
| | <p>(ii) Antacids are the medicines used to control acidity in stomach. Ex – mixture of aluminium and magnesium hydroxide / sodium hydrogen carbonate / Zantac / Ranitidine (or any other suitable example)</p> <p>(iii) No, Excessive antacid can make the stomach alkaline and trigger the production of more acid.</p> | <p>1+ ½</p> <p>½ + 1</p> |
| 24 | <p>a) $E_{\text{cell}} = E_{\text{cell}}^0 - \frac{0.0591 \text{ V}}{n} \log \frac{[\text{Al}^{3+}]^2}{[\text{Cu}^{2+}]^3}$</p> <p>$E_{\text{cell}}^0 = E_{\text{cell}} + \frac{0.0591 \text{ V}}{n} \log \frac{[\text{Al}^{3+}]^2}{[\text{Cu}^{2+}]^3}$</p> <p>$E_{\text{cell}}^0 = 1.98 \text{ V} + \frac{0.0591 \text{ V}}{6} \log \frac{(0.01)^2}{(0.01)^3}$</p> <p>$E_{\text{cell}}^0 = 1.98 \text{ V} + \frac{0.0591 \text{ V}}{6} \log 10^2$</p> <p>$E_{\text{cell}}^0 = 1.98 \text{ V} + \frac{0.0591 \text{ V}}{6} \times 2 \times \log 10 \quad [\because \log 10 = 1]$</p> <p>$E_{\text{cell}}^0 = 1.98 \text{ V} + \frac{0.0591 \text{ V}}{6} \times 2$</p> <p>$E_{\text{cell}}^0 = 1.98 \text{ V} + 0.0197 \text{ V}$</p> <p>$E_{\text{cell}}^0 = 1.9997 \text{ V}$</p> <p>(b) A , because its E^0 value is more negative.</p> | <p>1</p> <p>1</p> <p>1</p> <p>1+1</p> |
| | OR | |
| 24 | <p>(a) $\Lambda_{\text{m}}^{\text{c}} = \kappa \times 1000 / C$</p> <p>$= 3.905 \times 10^{-5} \times 1000 / 0.001$</p> <p>$= 39.05 \text{ S cm}^2/\text{mol}$</p> <p>$\text{CH}_3 \text{COOH} \rightarrow \text{CH}_3\text{COO}^- + \text{H}^+$</p> <p>$\Lambda^{\circ} \text{CH}_3\text{COOH} = \lambda^{\circ} \text{CH}_3 \text{COO}^- + \lambda^{\circ} \text{H}^+$</p> <p>$= 40.9 + 349.6$</p> <p>$\Lambda^{\circ} \text{CH}_3\text{COOH} = 390.5 \text{ S cm}^2/\text{mol}$</p> | <p>½</p> <p>1</p> |

| | | |
|----|---|---|
| | $\alpha = \frac{A_m}{A_m^0}$ $= 39.05 / 390.5$ $= 0.1$ <p>(b) Device used for the production of electricity from energy released during spontaneous chemical reaction and the use of electrical energy to bring about a chemical change. The reaction gets reversed / It starts acting as an electrolytic cell & vice – versa.</p> | $\frac{1}{2}$ 1 1 1 |
| 25 | <p>(a)</p> <p>i) Ability of oxygen to form multiple bond with Mn metal.</p> <p>ii) Cr^{2+} is oxidized to Cr^{3+} which has stable d^3 / t_{2g}^3 orbital configuration</p> <p>iii) Cu^{2+} has unpaired electron while Zn^{2+} has no unpaired electron.</p> <p>(b)</p> <p>i) $2\text{MnO}_2 + 4\text{KOH} + \text{O}_2 \xrightarrow{\triangle} 2\text{K}_2\text{MnO}_4 + 2\text{H}_2\text{O}$</p> <p>ii) $\text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ + 6\text{I}^- \longrightarrow 2\text{Cr}^{3+} + 7\text{H}_2\text{O} + 3\text{I}_2$ (balanced equation is required)</p> | 1 1 1 1 1 |
| | OR | |
| 25 | <p>i) Mn. It has maximum unpaired electrons.</p> <p>ii) Cr</p> <p>iii) Sc</p> <p>iv) Manganese. Mn^{3+} to Mn^{2+} results in the stable half filled (d^5) configuration.</p> | $\frac{1}{2} + 1$ 1 1 $\frac{1}{2} + 1$ |
| 26 | <p>(a)</p> <p>(i) A: CH_3CHO , B: $\text{CH}_3\text{CH}=\text{N}-\text{OH}$</p> <p>(ii) A: CH_3COOH , B: CH_3COCl</p> <p>(b)</p> <p>(i) Heat both compounds with NaOH and I_2, $\text{C}_6\text{H}_5\text{COCH}_3$ forms yellow ppt of CHI_3 whereas $\text{C}_6\text{H}_5\text{CHO}$ does not.</p> <p>(ii) Add ammoniacal solution of silver nitrate (Tollen's reagent) to both the compounds, HCOOH gives silver mirror but CH_3COOH does not. (or any other suitable test)</p> <p>(C) $\text{CH}_3\text{CHO} < \text{CH}_3\text{CH}_2\text{OH} < \text{CH}_3\text{COOH}$</p> | $\frac{1}{2} + \frac{1}{2}$ $\frac{1}{2} + \frac{1}{2}$ 1 1 1 |

| | | |
|----|--|---|
| | OR | |
| 26 | <p>(a)</p> $\text{>C=O} \xrightarrow[-\text{H}_2\text{O}]{\text{NH}_2\text{NH}_2} \text{>C=NH}_2 \xrightarrow[\text{heat}]{\text{KOH/ethylene glycol}} \text{>CH}_2 + \text{N}_2$ <p>(b) $\text{C}_6\text{H}_5\text{COCH}_3 < \text{CH}_3\text{COCH}_3 < \text{CH}_3\text{CHO}$</p> <p>(c) Because of resonance in carboxylic group the carbonyl group loses a double bond character.</p> <p>(d) $\text{CH}_3\text{CH}_2\text{CH}=\text{CH}-\text{CH}_2\text{CHO}$</p> <p>(e) A : $\text{CH}_3\text{CH}_2\text{CHO}$ B : CH_3COCH_3</p> | <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>$\frac{1}{2} + \frac{1}{2}$</p> |

| Name | Signature | Name | Signature |
|-------------------------------|-----------|---------------------------|-----------|
| Dr. (Mrs.) Sangeeta Bhatia | | Sh. S.K. Munjal | |
| Dr. K.N. Uppadhya | | Sh. D.A. Mishra | |
| Prof. R.D. Shukla | | Sh. Rakesh Dhawan | |
| Dr. (Mrs.) Sunita Ramrakhiani | | Ms. Nirmala Venkateswaran | |
| Sh. S. Vallabhan, Principal | | Mrs. Deepika Arora | |
| Mr. K.M. Abdul Raheem | | Ms. Minakshi Gupta | |
| Mrs. Sushma Sachdeva | | Sh. Mukesh Kaushik | |
| Ms. Seema Bhatnagar | | Mr. Roop Narayan | |
| Sh. Pawan Singh Meena | | Ms. Garima Bhutani | |
| Sh. Praveen Kumar Agrawal | | | |